

COLONOSCOPY PREPARATION WITH MOVIPREP			
PLEASE PURCHASE	<ul style="list-style-type: none"> • MOVIPREP Kit (by prescription) • The kit contains 2 Pouch A and 2 Pouch B packets 		
7 DAYS BEFORE	<ul style="list-style-type: none"> • Stop fiber supplements and iron supplements, including multivitamins 		
3 DAYS BEFORE	<table border="0"> <tr> <td style="vertical-align: top;"> <p>Begin a LOW FIBER Diet including:</p> <ul style="list-style-type: none"> • White or sour dough bread • White Rice, Pasta • Eggs, Chicken, Turkey, Fish • Cheese, Yogurt, Milk, Ice Cream, Pudding • Canned or cooked fruit/vegetables without skin or peel (carrots, cucumbers, potatoes) • Pureed fruits/vegetables (smoothies, strained puree soups) • Creamy peanut or almond butter </td> <td style="vertical-align: top;"> <p>High Fiber Foods to AVOID:</p> <ul style="list-style-type: none"> • Raw fruits/vegetables with seeds, skin, membranes • Cooked/canned fruit/vegetables with skin/seeds/peel • Whole wheat or multigrain bread or crackers • Nuts, Seeds • Popcorn • Oatmeal, bran cereal • Brown rice, quinoa </td> </tr> </table>	<p>Begin a LOW FIBER Diet including:</p> <ul style="list-style-type: none"> • White or sour dough bread • White Rice, Pasta • Eggs, Chicken, Turkey, Fish • Cheese, Yogurt, Milk, Ice Cream, Pudding • Canned or cooked fruit/vegetables without skin or peel (carrots, cucumbers, potatoes) • Pureed fruits/vegetables (smoothies, strained puree soups) • Creamy peanut or almond butter 	<p>High Fiber Foods to AVOID:</p> <ul style="list-style-type: none"> • Raw fruits/vegetables with seeds, skin, membranes • Cooked/canned fruit/vegetables with skin/seeds/peel • Whole wheat or multigrain bread or crackers • Nuts, Seeds • Popcorn • Oatmeal, bran cereal • Brown rice, quinoa
<p>Begin a LOW FIBER Diet including:</p> <ul style="list-style-type: none"> • White or sour dough bread • White Rice, Pasta • Eggs, Chicken, Turkey, Fish • Cheese, Yogurt, Milk, Ice Cream, Pudding • Canned or cooked fruit/vegetables without skin or peel (carrots, cucumbers, potatoes) • Pureed fruits/vegetables (smoothies, strained puree soups) • Creamy peanut or almond butter 	<p>High Fiber Foods to AVOID:</p> <ul style="list-style-type: none"> • Raw fruits/vegetables with seeds, skin, membranes • Cooked/canned fruit/vegetables with skin/seeds/peel • Whole wheat or multigrain bread or crackers • Nuts, Seeds • Popcorn • Oatmeal, bran cereal • Brown rice, quinoa 		
2 DAYS BEFORE	<ul style="list-style-type: none"> • Continue LOW FIBER Diet • Avoid solid foods after midnight 		
1 DAY BEFORE	<ul style="list-style-type: none"> • MORNING: Start a CLEAR LIQUID Diet <ul style="list-style-type: none"> ○ Water ○ Apple, White Grape, White Cranberry Juices without Pulp ○ Clear Soup/Broth ○ Tea or Black coffee (no milk/creamer) ○ Gatorade/Powerade (no red, purple or orange color) ○ Jello (no red, purple, or orange color) ○ Popsicles (no red, purple, or orange color) • 6PM: Take MOVIPREP Dose 1 <ul style="list-style-type: none"> ○ Empty one POUCH A and one POUCH B into the container ○ Add cold water to the top of the line and dissolve ○ Drink solution down to the next mark every 15 minutes ○ Drink 16 oz of additional clear liquid after finishing the Moviprep • Bowel movements should begin 1-2 hours after starting MOVIPREP but effect can sometimes be delayed by up to 4-6 hours 		

DAY OF COLONOSCOPY

5 HOURS BEFORE	<ul style="list-style-type: none"> • Take MOVIPREP Dose 2 <ul style="list-style-type: none"> ○ Follow the same instructions as outlined for Dose 1 • By the end of the cleanse your stool should be a clear liquid with yellow or green tinge
3 HOURS BEFORE	<ul style="list-style-type: none"> • Stop drinking/eating clear liquids • Do not chew gum or mints • Take regular medications with a small sip of water the morning of your procedure unless otherwise instructed by your physician
30 MIN BEFORE	<ul style="list-style-type: none"> • Arrive at endoscopy suite 30 MIN PRE PROCEDURE (no later!) • Bring list of current medications including doses • Bring a Photo ID and Insurance Card • Bring any applicable Referrals • Please do not bring any valuables

GENERAL INSTRUCTIONS:

IF YOU HAVE DIABETES	<ul style="list-style-type: none"> • Contact your primary care doctor or endocrinologist for instructions regarding your medications
IF YOU TAKE MEDICATION TO THIN YOUR BLOOD	<ul style="list-style-type: none"> • Contact your gastroenterologist for further instructions at (212) 427-8761 • Medications include but are not limited to: <ul style="list-style-type: none"> ○ Heparin, Lovenox, Warfarin (Coumadin), Rivaroxaban (Xarelto), Apixaban (Eliquis), Edoxaban (Lixiana, Savaysa), Betrixaban (Bevyxxa), Clopidigrel (Plavix), Prasugrel (Effient), Brilinta (Ticagrelor)
IF YOU TAKE ASPIRIN OR BLOOD PRESSURE MEDICATIONS	<ul style="list-style-type: none"> • You may continue to take these medications
ESCORT	<ul style="list-style-type: none"> • Please ensure that you have an escort (responsible adult over the age of 18) to pick you up at the end of the procedure.