MANHATTAN GASTROENTEROLOGY

147 East 26th Street
New York, NY 10010
(212) 427-8761
Manhattangastroenterology.com

2-DAY COLONOSCOPY PREPARATION: MIRALAX EXTENDED		
PLEASE PURCHASE	Over the counter at local pharmacy • **Dulcolax (bisacodyl) 5 mg pills** (need 3 pills total) • **Miralax 238 gram bottle x 2** • **Gatorade: 28 (or 32) oz bottles x3** (No Red, Purple, or Orange Color)	
7 DAYS BEFORE	• **Stop any GLP-1 medications** (injection or oral) • **Stop fiber supplements and iron supplements** (including multivitamins) • Cease smoking any marijuana, tobacco, or nicotine products	
4 DAYS BEFORE	**Begin a LOW FIBER Diet including:** • White or sourdough bread • White Rice, Pasta • Eggs, Chicken, Turkey, Fish • Cheese, Yogurt, Milk, Ice Cream, Pudding • Canned or cooked fruit/vegetables without skin or peel (carrots, cucumbers, potatoes) • Pureed fruits/vegetables (smoothies, strained puree soups) • Creamy peanut or almond butter	**High Fiber Foods to AVOID:** • Raw fruits/vegetables with seeds, skin, membranes • Cooked/canned fruit/vegetables with skin/seeds/peel • Whole wheat or multigrain bread or crackers • Nuts, Seeds • Popcorn • Oatmeal, bran cereal • Brown rice, quinoa

2 DAYS BEFORE	• **Continue LOW FIBER Diet**
	• **7 PM:**
	  ○ Pour 7 capfuls of MiraLAX (half of a 14-dose, 238 g bottle) into a bottle of Gatorade. Shake until completely dissolved.
	  ○ Drink the first bottle, one 8 oz glass every 15 minutes, until finished.	

1 DAY BEFORE	• **MORNING: Start a CLEAR LIQUID diet**
	  ○ Apple, White Grape, White Cranberry Juices without Pulp
	  ○ Clear Soup/Broth
	  ○ Tea or Black coffee (no milk/creamer)
	  ○ Gatorade/Powerade (no red, purple, or orange color)
	  ○ Jell-O (no red, purple, or orange color)
	  ○ Popsicles (no red, purple, or orange color)
	• **6 PM:** Take Dulcolax 3 tablets with sips of water
	• **7 PM:**
	  ○ Pour 7 capfuls of MiraLAX (half of a 14-dose, 238 g bottle) into each of the next two bottles of Gatorade. Shake until completely dissolved.
	  ○ Drink the second bottle, one 8 oz glass every 15 minutes, until finished.
	  ○ Refrigerate the third bottle for the next part of your prep.
	• Bowel movements should begin 1-2 hours after starting the Miralax/Gatorade mix, but the effect can be delayed by up to 4-6 hours	

DAY OF COLONOSCOPY	
5 HOURS BEFORE	• Drink the third bottle of Miralax/Gatorade mix ○ one 8 oz glass every 15 minutes, until finished. • By the end of the cleanse, your stool should be a clear liquid with a yellow or green tinge
3 HOURS BEFORE	• Stop drinking/eating clear liquids • Do not chew gum or mints • Take regular medications with a small sip of water the morning of your procedure unless otherwise instructed by your physician
30 MIN BEFORE	• **Arrive at endoscopy suite 30 MIN PRE PROCEDURE (no later)** • **If you are a premenopausal female, you will need to submit a urine sample upon office check-in** • Bring list of current medications including doses • Bring a Photo ID and Insurance Card • Bring any applicable Referrals • Please do not bring any valuables

GENERAL INSTRUCTIONS AND TIPS

IF YOU HAVE DIABETES	• Contact your primary care doctor or endocrinologist for instructions regarding your medications
IF YOU TAKE MEDICATION TO THIN YOUR BLOOD	• Contact your gastroenterologist for further instructions at (212) 427-8761 • Medications include but are not limited to: ○ Heparin, Lovenox, Warfarin (Coumadin), Rivaroxaban (Xarelto), Apixaban (Eliquis), Edoxaban (Lixiana, Savaysa), Betrixaban (Bevyxxa), Clopidogrel (Plavix), Prasugrel (Effient), Brilinta (Ticagrelor)
IF YOU TAKE ASPIRIN OR BLOOD PRESSURE MEDICATIONS	• You may continue to take these medications
ESCORT	• Please ensure that you have an escort (responsible adult over the age of 18) to pick you up at the end of the procedure